

Removal of Impacted Canines

Department of Oral and Maxillofacial Surgery

Introduction

The canines, or eye teeth, normally erupt into the mouth between the ages of 11 and 13. Sometimes, one or both canines develop in the wrong position and become impacted. The majority impact towards the roof of the mouth (palate) behind the front teeth but some impact on the cheek side (buccal).

What is meant by impacted?

A tooth that is prevented from erupting into its normal functional position by bone, tooth or soft tissue.

How does it happen?

Upper canines are the last teeth to come through and are therefore more likely to become impacted. It can be caused by missing lateral incisors which provide lack of guidance to the canine, lack of space for eruption due to crowding and in some cases there is a family history.

Diagnosis

Diagnosis is by clinical and radiographic (x-ray) examination. Radiographic examination usually involves taking 2 radiographs (x-rays).

Treatment Options

- Leave the impacted canine as it is and accept the appearance. There is a small chance that the canine could cause problems in the future by damaging the neighbouring teeth or by formation of a cyst.
- Surgical removal of the impacted tooth. This involves a short operation, often under general anaesthetic. This is done as a day case procedure meaning that you can go home that day. The gum needs to be lifted up so that the tooth can be identified and removed. Afterwards there will be some stitches between the teeth. The mouth will be a little uncomfortable for a few days up to a week. Usually normal painkillers such as ibuprofen and / or Paracetamol are sufficient.
- Exposure & bonding of the tooth to bring it into the correct position. This may not be possible if the tooth is in a difficult position.

Your surgeon and/or orthodontist will advise you on the best treatment option.

Additional information can also be obtained by logging on to <http://www.baoms.org.uk.sitemap.asp?id=20>

Other web links

National Institute of Dental and Craniofacial Research

Patient.co.uk

Department of Oral and Maxillofacial Surgery
St. Peter's Hospital Telephone: 01932 872000 ext. 2493
Ashford Hospital Telephone 01784 884009

Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty. If they cannot resolve your concern, please contact our Patient Advice and Liaison Service (PALS) on 01932 723553 or email pals@asph.nhs.uk. If you still remain concerned please contact our Complaints Manager on 01932 722612 or email complaints@asph.nhs.uk.

Author: M Dave

Department: Oral and Maxillofacial Surgery

Version: 2

Published: Nov 2008

Review: Nov 2011

We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.



To use the Text Relay service, prefix all numbers with 18802.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید.
ਜੇ ਤੁਹਾਨੂੰ ਭਰਜਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ ਫੋਨ ਕਰੋ: 01932 723553
اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں
Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553
यदि आपको अनुवाद की जरूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553
Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Ashford Hospital London Road, Ashford, Middlesex, TW15 3AA Tel: **01784 884488**
St. Peter's Hospital Guildford Road, Chertsey, Surrey, KT16 0PZ Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk