

Review

Orofacial pain

Kenneth M. Hargreaves*

*Departments of Endodontics, Pharmacology, Physiology and Surgery, University of Texas Health Science Center at San Antonio, San Antonio, TX 78229, USA**Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.***1. Introduction**

Pain in the oral and craniofacial system represents a major medical and social problem. Indeed, a U.S. Surgeon General's report on orofacial health concludes that, "...oral health means much more than healthy teeth. It means being free of chronic oral-facial pain conditions..." [172]. Community-based surveys indicate that many subjects commonly report pain in the orofacial region, with estimates of >39 million, or 22% of Americans older than 18 years of age, in the United States alone [108]. Other population-based surveys conducted in the United Kingdom [111,112], Germany [91], or regional pain care centers in the United States [54] report similar occurrence rates [135]. Importantly, chronic widespread body pain, patient sex and age, and psychosocial factors appear to serve as risk factors for chronic orofacial pain [1,2,92,99,138]. In addition to its high degree of prevalence, the reported intensities of various orofacial pain conditions are similar to that observed with many spinal pain disorders (Fig. 1). Moreover, orofacial pain is derived from many unique target tissues, such as the meninges, cornea, tooth pulp, oral/nasal mucosa, and temporomandibular joint (Fig. 2), and thus has several unique physiologic characteristics compared with the spinal nociceptive system [23]. Given these considerations, it is not surprising that accurate diagnosis and effective management of orofacial pain conditions represents a significant health care problem.

Publications in the field of orofacial pain demonstrate a steady increase over the last several decades (Fig. 3). This is a complex literature; a recent bibliometric analysis of orofacial pain articles published in 2004–2005 indicated that 975 articles on orofacial pain were published in 275 journals from authors representing 54 countries [142]. Thus, orofacial pain disorders represent a complex constellation of conditions with an equally diverse literature base. Accordingly, this review will focus on a summary of major research foci on orofacial pain without attempting to provide a comprehensive review of the entire literature.

2. Physiologic studies on trigeminal pain

Several reviews are available that document the historical development of physiologic research on the trigeminal nociceptive

system [33,53,114,120,149]. More recent studies have characterized differences in electrophysiological [82], anatomical [10], or pharmacological [42,80] properties of trigeminal afferents innervating distinct target tissues. Collectively, many of these studies provide support for the hypothesis that target tissue interactions with trigeminal neuron terminals, via either soluble factors such as neurotrophins [48], or by integrin binding to extracellular matrix molecules [22,25], regulate the expression or trafficking of neuronal proteins, including ion channels and receptors [90,139] or second messenger signaling pathways [25]. Thus, the presence of unique target tissues innervated by trigeminal afferent fibers likely contributes to differences in the responsiveness of these neurons. A recent review characterizes differences between the trigeminal and spinal afferent systems under basal conditions [23]. Table 1 illustrates differences between the trigeminal and spinal systems after various forms of injury. Collectively, these studies indicate that the trigeminal system has many unique features that may contribute to distinct response patterns to tissue injury.

The hypothesis of peripheral regulation of neuronal phenotype has been expanded by the recognition that estradiol selectively alters gene transcription in trigeminal neurons with increased expression of neuropeptides, such as prolactin, that are capable of sensitizing neuronal responses to capsaicin or noxious heat [49]. Additional studies have demonstrated that trigeminal peptidergic neurons undergo morphological changes ("sprouting") in response to injury-induced inflammation in target tissues [33]. In contrast, there is a lack of sympathetic fiber sprouting in trigeminal ganglion cells, unlike the well-recognized occurrence in the spinal system [19,29,63]. Thus, an emerging body of evidence reveals the dynamic and specific responsiveness of the trigeminal system to either injury of its various target tissues or to the presence of certain gonadal steroids.

Other studies have employed cultured trigeminal ganglia (TG) to evaluate cellular mechanisms of neuronal activation. For example, cannabinoids activate a calcineurin pathway leading to the rapid dephosphorylation and desensitization of transient receptor potential cation channel, subfamily V, member 1 (TRPV1), thereby contributing to an ionotropic mechanism for peripheral cannabinoid antinociception [4,5,89,133]. Moreover, accumulating evidence indicates a functional cross-desensitization between TRPV1 and TRP subfamily A, member 1 (TRPA1) on trigeminal neurons, possibly via formation of a heteromer [4,6,146,147]. Additional studies have used cultured TG to demonstrate that opioid receptors are

* Address: Department of Endodontics, UTHSCSA, 7703 Floyd Curl Drive, San Antonio, TX 78229, USA. Tel.: +1 210 567 3388; fax: +1 210 567 3389.

E-mail address: Hargreaves@uthscsa.edu




Fig. 1. Comparison of pain intensity among spinal and orofacial pain disorders using the McGill Total Rank Pain Index (PRI(T)). The PRI(T) is an ordinal scale consisting of the sum of the ranks of words in each of the 20 sub-categories on the McGill Pain Questionnaire and ranges from 0 to 78. Data taken from: [14,35,71,73,116,117,150,164,176,183,184].




Fig. 2. Unique target tissues innervated by the trigeminal sensory system. Taken from Bereiter et al. [23], with permission.

expressed on sensory neurons but are not coupled to inhibitory signaling pathways under basal experimental conditions. Instead, pretreatment with arachidonic acid or with agonists to receptors coupled to $G\alpha_q$ signaling pathways (eg, bradykinin, trypsin) is required to induce the rapid development of a functional competence for opioid receptor signaling to $G\alpha_i$ pathways leading to inhibition of neuronal activities [24,25,60,131,132]. These cellular findings are consistent with the observation that opioids have little efficacy for peripheral antinociception under basal conditions, but rapidly gain functional competence following injection of inflammatory mediators [145] or the development of inflammation.

Recent studies have employed cultured trigeminal neurons to identify endogenous TRPV1 agonists [130,134]. Heating of isolated superfused peripheral tissue to a noxious temperature range leads to the release of oxidized linoleic acid metabolites (OLAMs), including 9- and 13-hydroxyoctadecadienoic acid (HODE). The administration of synthetic 9- and 13-HODE (and their oxoODE metabolites) selectively activates TRPV1, leading to inward currents, increased accumulation of intracellular calcium, and triggering exocytosis of neuropeptides from TG neurons and thermal allodynia. These effects are blocked by TRPV1 antagonists and are

observed only in trigeminal neurons from wild-type mice but not TRPV1 knockouts [130]. Moreover, the intracellular delivery of compounds that block OLAM formation (eg, nordihydroguaiaretic acid) or a combination of anti-9- and anti-13-HODE antibodies both significantly inhibit heat-evoked activation of trigeminal neurons. Collectively, these findings strongly implicate the OLAMs as a family of endogenous TRPV1 agonists. Interestingly, the pronounced effect of TRPV1 antagonists for blocking heat hyperalgesia in inflammation as well as mechanical allodynia (after intrathecal administration) has led to the hypothesis that an endogenous TRPV1 system might be activated under conditions of tissue injury. In support of this hypothesis, the administration of anti-OLAM antibodies produces a peripherally mediated thermal antinociception and a centrally mediated blockade of mechanical allodynia in the complete Freund's model of inflammation [130,134]. Thus, the OLAM system appears to contribute to acute heat detection by TRPV1 and to regulate more persistent conditions such as inflammatory pain.

Future research directions may include preclinical studies focusing on mechanisms underlying differences between trigeminal and spinal pain conditions, mechanisms of sex-dependent differences in pain transduction and processing, and on the biological basis and pharmacological regulation of acute and chronic orofacial pain conditions. Many of these studies would be promoted by the development of standardized preclinical pain models and assessment methods.

In addition to research on the biological mechanisms of nociceptive transmission, numerous clinical studies have described strong psychosocial/disability components to orofacial pain. Indeed, some diagnostic classification schemes differentiate the dimension of tissue contributions from psychosocial/disability factors [55] contributing to orofacial pain disorders. These studies demonstrate that the orofacial pain patient is confronted with a complex, multidimensional disorder that is best managed with appropriate treatment for all underlying factors [2,46,56,61,85,97,154].

3. Studies on trigeminal inflammatory disorders

Many translational studies have evaluated acute inflammatory injury to the trigeminal system. The dental impaction pain model has been developed as a standard clinical method for evaluating many analgesic drugs [43,45,115]. Other investigators have used this model of acute inflammatory pain to evaluate preemptive anesthesia [66,68], activation of endogenous opioid analgesic systems [70,78,104,105], local release of inflammatory mediators as collected by implanted microdialysis probes [50,67,77,162], other physiologic mechanisms [76], or the association of genetic polymorphisms with postoperative pain [95,96,98]. This clinical model has several notable advantages, including participation of relatively healthy subjects not taking concurrent drugs, standardized surgical procedures leading to reduced variance, and relatively large numbers of potential participants. Collectively, these studies on patients undergoing surgical dental extractions have contributed greatly to evaluation of analgesics, anesthetics, and anxiolytics, as well as basic biological research on human subjects.

Other studies have focused on chronic inflammation of the oral and craniofacial region. Clinical studies on irreversible pulpitis in teeth ("toothache") have demonstrated that this condition of bacterial-induced inflammation/necrosis is associated with significant changes in expression of ion channels [8,16,83,110,178,182], receptors [106], and neuropeptides [32,33]. Moreover, inflammation of a single tooth in patients appears sufficient to trigger central sensitization [59,94,127,151]. Animal studies on inflammation in the trigeminal region have demonstrated target-site-dependent differences in sensitization/activation [62,79,175] as well as sex-dependent differences in neuronal activities [20,126,168].




Fig. 3. Rates of articles published on orofacial pain. Data were acquired from a PubMed search (August 2010) using the search criteria of: (orofacial or trigeminal or temporomandibular or dental or tooth) and (pain or headache or hyperalgesia or allodynia or nociceptor or nociceptive).

Future research directions on trigeminal inflammatory disorders may include preclinical and translational clinical studies focusing on mechanisms underlying the development and maintenance of inflammatory hyperalgesia/allodynia. Importantly, the clinical condition of pulpitis results in a very restricted pain locus (often within a tissue volume of <200 μ L), intense pain reports [73], and dynamic neuronal and immunoplasticity. Thus, the pulpitis pain model is important not only from the perspective of high prevalence [108] and health care disparity [174], but also serves as a useful model for translational research [27].

4. Studies on trigeminal neuropathic disorders

The orofacial region has unique neuropathic pain disorders not seen in the spinal system, including trigeminal neuralgia and glossopharyngeal neuralgia [15]. Numerous clinical reports document these and other orofacial neuropathic or neuritic/neuralgic pain conditions and their responsiveness to surgical or pharmacological treatments [15,38,118,124,129,168,184]. Several etiologic factors appear to contribute to the development of neuropathic pain disorders. Proposed mechanisms include injury/compression to the trigeminal nerve, inflammatory insult (possibly including glial contributions), or infection with herpes virus [3,15,17,40,124,125,177]. However, not all injuries to the trigeminal nerve lead to neuropathic pain disorders; indeed, the incidence of neuropathic pain after injury to orofacial structures is relatively low after dental treatment [34,109,137], facial trauma [17], orthognathic surgery [39], tooth extraction [26,36,143,173], or placement of dental implants [72]. This apparent resistance of the trigeminal system to development of neuropathic conditions is an interesting clinical observation that should prompt preclinical research comparing trigeminal to spinal afferent systems for susceptibility to neuropathic pain disorders. It is interesting that the trigeminal system appears programmed for periodic loss of innervated structures during postnatal development, with the shedding of 20 deciduous teeth per person, with minimal development of neuropathic pain conditions.

Several risk factors for trigeminal neuralgia have been found, including multiple sclerosis [44,140] and hypertension [93]. Additional studies have reported changes in the expression of ion channels (eg, NaV1.3, 1.7, 1.8, TRPA1) in surgical biopsies collected from patients suffering from neuropathic orofacial pain [119,153]. Animal models of trigeminal neuropathic pain have been developed and include chronic constriction injury of the infraorbital nerve as well as transection of the inferior alveolar nerve [3,177]. Interestingly, both preclinical and clinical studies have implicated constriction of peripheral nerves as an etiologic mechanism for inducing neuropathic pain via alteration in primary afferent functions [81], although certain cortical changes have been reported as well [28]. This has led to the development of clinical surgical decompression procedures to treat patients with trigeminal neu-

ralgia [166,167]. Several preclinical studies have implicated ion channels and endothelin receptors as well as glial mechanisms in contributing to the development of these models of neuropathic pain conditions [11,40,81,125].

5. Studies on chronic trigeminal myofascial and joint pain

The diagnosis and management of many chronic orofacial pain conditions has been greatly hampered by confusion in determining etiologies from the temporomandibular joint versus myofascial sources. This has led to clinical studies difficult to interpret and diagnostic classifications that did not have a strong biological basis due to the lack of differentiation between joint and muscle contributions to the patient's pain condition. Clinical studies on myofascial pain or temporomandibular dysfunction (TMD) were considerably improved by the development of the Research Diagnostic Criteria [55,56], which highlighted the need for developing standardized diagnostic methods and definitions. Considerable evidence has been published demonstrating that patient sex/gender and exposure to sex steroids serve as risk factors for developing chronic orofacial pain conditions [61,65,100–103]. However, this is not observed in all studies, and other risk factors such as chronic widespread body pain, a prior history of physical abuse, or health anxiety have also been reported to be associated with the development of chronic orofacial pain disorders [1,55,61,92,100,107]. The reasons why some but not all studies detect sex/gender as a significant risk factor for orofacial pain disorders is not clear, but may be due to differences in patient populations, case definitions, or experimental approaches. Related preclinical studies have demonstrated that trigeminal neurons express estrogen receptors and undergo dramatic changes in gene expression [9,21,49] or firing rates [62] following exposure to estradiol.

Other clinical studies have focused on synovial fluid levels of inflammatory mediators to test for other possible biological mechanisms [31,160] or have evaluated the role of peripheral glutamate receptors in triggering myofascial pain [7,12]. A very interesting approach is the application of genetics to patients with TMD. A haplotype of the catechol-O-methyltransferase gene in patients is associated with reduced responsiveness to experimental pain and to reduced risk for TMD [47]. Moreover, a mechanistic hypothesis for the protective effect of this haplotype has been advanced [121], and TMD patients with this catechol-O-methyltransferase haplotype respond with increased analgesia from drugs such as propranolol [169].

6. Studies on other orofacial pain conditions

Many other orofacial pain disorders also have been evaluated. The trigeminal autonomic cephalgias include cluster headache,

Table 1
Comparison of the trigeminal and spinal afferent systems after injury.

Marker	Injury model	Comparison	Authors
Galanin	Axotomy	TG ~ DRG for upregulation	Arvidsson et al. [13]; Zhang et al. [185]
NPY	Axotomy	TG ~ DRG for upregulation	Arvidsson et al. [13]; Zhang et al. [185]
Sympathetic fiber sprouting into ganglion and basket formation	Axotomy or CCI	TG: No DRG: Yes	Bongehiellm et al. [29]; Benoliel et al. [19]
Sympathetic fiber sprouting into ganglion	NGF infusion icv X 14d	DRG > TG	Nauta et al. [122]
SNS/PN3 = NaV1.8	Axotomy	TG: Downregulation followed by normalization DRG: Persistent downregulation	Bongehiellm et al. [30]
Ankyrin(G)	Axotomy	TG: Persistent downregulation	Bongehiellm et al. [30]
Ectopic firing of afferents	Axotomy	TG < DRG	Tal and Devor [165]
Augmented excitability	Axotomy	TG ~ DRG	Tal and Devor [165]; Zhang et al. [186]; Cherkas et al. [37]
Frequency and rhythmicity of spontaneous discharges	Tight ligation of infraorbital vs sciatic nerves	DRG had significantly greater spontaneous discharge rate than TG neurons for both myelinated and unmyelinated fibers. DRG afferents had rhythmic discharge rates (not seen with TG)	Tal and Devor [165]
Satellite glial cells	Axotomy	TG ~ DRG for upregulation of GFP, proliferation	Woodham et al. [181]; Stephenson et al. [158]; Cherkas et al. [37]
NOS	Axotomy	TG ~ DRG for upregulation	Hokfelt et al. [84]
P2X3 & ATF-3 expression	Partial axotomy	TG ~ DRG	Tsuzuki et al. [171]
GM3 ganglioside.	Knockout GM2/GD2 and the GD3 synthase gene	Facial wounding > Rest of the body	Inoue et al. [87]
Peripheral chromatolysis	LiCl	With peripheral nerve degeneration	Levine et al. (2004)
Sensory neuropathy with neuronal degeneration	Sjogren's Syndrome	TG ~ DRG	Malinow et al. [113]
Infectivity of contralateral ganglia	Herpes simplex virus-1 (HSV) infection	70% of TG contralateral to side of HSV injection produced infections after inoculation, whereas only 10% of contralateral DRG produced infections.	Thackray et al. [170]
HSV polypeptide ICP4 (VP175) expression in ganglia	Herpes simplex virus-1 (HSV) infection	TG ~ DRG	Pepose et al. [136]
Viral replication and degradation of host cells, mRNA	Herpes simplex virus-1 (HSV) infection	TG ~ DRG with wild-type HSV more virulent in both ganglia than HSV mutants lacking virion host shutoff (vhs) protein	Smith et al. [156]
Infectivity of ganglia	Simian varicella virus (SVV)	TG ~ DRG	White et al. [179]
Substance P in ganglia	Streptozotocin-diabetes	TG had 26% reduction ($P < 0.01$), but DRG = 11% non-significant reduction	Robinson et al. [144]
Substance P in ganglia	mf rat (mutilated foot; an autosomal recessive sensory neuropathy with reduced pain responsiveness)	DRG < TG	Scaravilli [148]
Caspase-3 mediated neuronal apoptosis	Knockout of Rb (retinoblastoma tumor suppressor protein)	TG ~ DRG for protection from apoptosis in double knockout of Rb and caspase-3 compared to single Rb knockout	Simpson et al. [152]
Number of neurons in ganglia	TRKA knockout	TG ~ DRG for extensive neuronal loss	Smeyne et al. [155]
Reactivation of virus	HSV mutant with gamma 34.5 gene deletion	TG > resistant to reactivation than DRG	Spivack et al. [157]
Wide-spread numbness and pain 4–12 d after antibiotic treatment	Acute sensory neuronopathy syndrome in humans	TG ~ DRG	Sterman et al. [159]
Pain	Trigeminal neuralgia in humans	TG: Yes (max/mand divisions > ophthalmic) DRG: No equivalent	Jannetta [88]; Sweet [161]; Wilkins [180]; Goya et al. [69]; Hamlyn [74,75]; Tacconi and Miles [163]
Spontaneous behavior	Formalin	OVX females exhibited significantly greater increase in formalin hyperalgesia after orofacial injection (upper lip) compared to hind paw injection. Result is consistent with hypothesis of a difference in sex steroid regulation of nociception between TG and DRG systems	Pajot et al. [128]

SNS, sensory neuron specific; PN3, peripheral nerve sodium channel type 3; NOS, nitric oxide synthase; P2X3, purinoceptor 3; ICP4, infected cell protein 4; TG, trigeminal ganglia; DRG, dorsal root ganglia; NPY, neuropeptide Y; CCI, chronic constriction injury; NGF, nerve growth factor; NOA, ATF-3, activating transcription factor 3; OVX, ovariectomized.

paroxysmal hemicrania, and unilateral neuralgiform headaches [41]. This collection of pain disorders is characterized by unilateral head pain in association with autonomic features such as tearing

and conjunctival involvement, and considerable research has shed light on pain referral patterns and issues related to proper diagnosis and treatment [18,51,58,64,86]. Most cases of trigeminal auto-

onomic cephalgia reflect primary headaches, although rare cases may be associated with pituitary tumors [41]. Pain is a major aspect of oral cancer [57] and often represents the initial symptom that prompts patients to seek health providers. Pain due to oral cancer may be due to soluble factors released from tumor cells, a localized inflammatory response to the tumor, or even nerve entrapment. Several recent studies have implicated the endothelin system and proteases (eg, PAR-2 receptor activation) in mediating mechanical allodynia experimental models of oral cancer pain [52,141]. Burning mouth syndrome is a rare disorder, commonly characterized by spontaneous burning pain and mechanical allodynia. Although idiopathic, it has many features of neuropathic pain and has been reported to be associated with altered peripheral expression of voltage-gated sodium channels [16].

7. Discussion

Orofacial pain disorders comprise a major and expensive component of health care and collectively have a high prevalence rate, a large range in pain intensity with a commensurate, often devastating impact on quality of life [149]. Although there are many common aspects of pain transduction and processing between the trigeminal and spinal systems, there are numerous examples of unique features in the peripheral and central components of the trigeminal pain system. Accordingly, ongoing basic and clinical research focused on acute and chronic orofacial pain conditions is required to understand the unique features of this pain system and to develop and evaluate better ways to treat patients with orofacial pain.

A major barrier for improved patient care and translational research is the lack of validated diagnostic criteria. Although efforts have been made to classify TMD patients with the Research Diagnostic Criteria for TMD, headache patients with the International Headache Society criteria, and orofacial pain with the American Academy of Orofacial Pain standards, clinical research indicates that each of these 3 methods is incomplete for comprehensive diagnosis of orofacial pain patients [18]. Thus, further research is critically required to establish a comprehensive, sensitive, and specific diagnostic classification scheme for all orofacial pain patients. This would provide a critical contribution to practitioners and foster the development of a powerful dataset for clinical research. In addition, recent studies have incorporated quality-of-life indices, which provide important additional information on clinical outcomes [123].

Taken together, orofacial pain conditions represent a highly prevalent spectrum of pain disorders with pain intensities similar to those observed with many chronic spinal pain conditions. However, the unique anatomical, biochemical, and associated psychosocial components provide compelling evidence for specific research focused on orofacial pain disorders.

Conflict of interest statement

The author denies any conflicts of interest.

Acknowledgement

This work was supported in part by R01 NS72890, R01 NS58655, R01 DA19585, and NCR R U54RR02438.

References

- [1] Aggarwal VR, Macfarlane GJ, Farragher TM, McBeth J. Risk factors for onset of chronic oro-facial pain—results of the North Cheshire oro-facial pain prospective population study. *Pain* 2010;149:354–9.
- [2] Aggarwal VR, Macfarlane TV, Macfarlane GJ. Why is pain more common amongst people living in areas of low socio-economic status? A population-based cross-sectional study. *Br Dent J* 2003;194:383–7 [discussion 380].
- [3] Ahn DK, Lim EJ, Kim BC, Yang GY, Lee MK, Ju JS, Han SR, Bae YC. Compression of the trigeminal ganglion produces prolonged nociceptive behavior in rats. *Eur J Pain* 2009;13:568–75.
- [4] Akopian AN, Ruparel NB, Jeske NA, Hargreaves KM. Transient receptor potential TRPA1 channel desensitization in sensory neurons is agonist dependent and regulated by TRPV1-directed internalization. *J Physiol* 2007;583:175–93.
- [5] Akopian AN, Ruparel NB, Jeske NA, Patwardhan A, Hargreaves KM. Role of ionotropic cannabinoid receptors in peripheral antinociception and antihyperalgesia. *Trends Pharmacol Sci* 2009;30:79–84.
- [6] Akopian AN, Ruparel NB, Patwardhan A, Hargreaves KM. Cannabinoids desensitize capsaicin and mustard oil responses in sensory neurons via TRPA1 activation. *J Neurosci* 2008;28:1064–75.
- [7] Alstergren P, Ernberg M, Nilsson M, Hajati AK, Sessle BJ, Kopp S. Glutamate-induced temporomandibular joint pain in healthy individuals is partially mediated by peripheral NMDA receptors. *J Orofac Pain* 2010;24:172–80.
- [8] Alvarado LT, Perry GM, Hargreaves KM, Henry MA. TRPM8 Axonal expression is decreased in painful human teeth with irreversible pulpitis and cold hyperalgesia. *J Endod* 2007;33:1167–71.
- [9] Amandusson A, Blomqvist A. Estrogen receptor-alpha expression in nociceptive-responsive neurons in the medullary dorsal horn of the female rat. *Eur J Pain* 2010;14:245–8.
- [10] Ambalavanar R, Moritani M, Dessem D. Trigeminal P2X3 receptor expression differs from dorsal root ganglion and is modulated by deep tissue inflammation. *Pain* 2005;117:280–91.
- [11] Amir R, Argoff CE, Bennett GJ, Cummins TR, Durieux ME, Gerner P, Gold MS, Porreca F, Strichartz GR. The role of sodium channels in chronic inflammatory and neuropathic pain. *J Pain* 2006;7:S1–S29.
- [12] Arendt-Nielsen L, Svensson P, Sessle BJ, Cairns BE, Wang K. Interactions between glutamate and capsaicin in inducing muscle pain and sensitization in humans. *Eur J Pain* 2008;12:661–70.
- [13] Arvidsson J, Raappana P, Diez M, Hokfelt T. Expression of neuropeptides in the rat mesencephalic trigeminal nucleus after peripheral axotomy. *Neuroreport* 1994;5:1269–72.
- [14] Barlas P, Craig JA, Robinson J, Walsh DM, Baxter GD, Allen JM. Managing delayed-onset muscle soreness: lack of effect of selected oral systemic analgesics. *Arch Phys Med Rehabil* 2000;81:966–72.
- [15] Baron R, Binder A, Wasner G. Neuropathic pain: diagnosis, pathophysiological mechanisms, and treatment. *Lancet Neurol* 2010;9:807–19.
- [16] Beneng K, Renton T, Yilmaz Z, Yiangou Y, Anand P. Sodium channel Na v 1.7 immunoreactivity in painful human dental pulp and burning mouth syndrome. *BMC Neurosci* 2010;11:71.
- [17] Benoliel R, Birenboim R, Regev E, Eliav E. Neurosensory changes in the infraorbital nerve following zygomatic fractures. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2005;99:657–65.
- [18] Benoliel R, Birman N, Eliav E, Sharav Y. The International Classification of Headache Disorders: accurate diagnosis of orofacial pain? *Cephalgia* 2008;28:752–62.
- [19] Benoliel R, Eliav E, Tal M. No sympathetic nerve sprouting in rat trigeminal ganglion following painful and non-painful infraorbital nerve neuropathy. *Neurosci Lett* 2001;297:151–4.
- [20] Bereiter DA, Benetti AP. Amino acid release at the spinomedullary junction after inflammation of the TMJ region in male and female rats. *Pain* 2006;126:175–83.
- [21] Bereiter DA, Cioffi JL, Bereiter DF. Oestrogen receptor-immunoreactive neurons in the trigeminal sensory system of male and cycling female rats. *Arch Oral Biol* 2005;50:971–9.
- [22] Bereiter DA, Cioffi JL, Bereiter DF, Zardeneta G, Milam SB. Local blockade of integrins in the temporomandibular joint region reduces Fos-positive neurons in trigeminal subnucleus caudalis of female rats produced by jaw movement. *Pain* 2006;125:65–73.
- [23] Bereiter DA, Hargreaves KM, Hu JW. Trigeminal mechanisms of nociception: peripheral and brainstem organization. In: Basbaum AI, Kaneko A, Shepherd GM, Westheimer G, Albright T, Masland RH, Dallos P, Oertel D, Firestein SJ, Beauchamp GK, Bushnell MC, Kaas JH, Gardner EP, editors. *The senses: a comprehensive reference*. San Diego: Academic Press; 2008. p. 435–60.
- [24] Berg KA, Patwardhan AM, Sanchez TA, Silva YM, Hargreaves KM, Clarke WP. Rapid modulation of micro-opioid receptor signaling in primary sensory neurons. *J Pharmacol Exp Ther* 2007;321:839–47.
- [25] Berg KA, Zardeneta G, Hargreaves KM, Clarke WP, Milam SB. Integrins regulate opioid receptor signaling in trigeminal ganglion neurons. *Neuroscience* 2007;144:889–97.
- [26] Berge TI. Incidence of chronic neuropathic pain subsequent to surgical removal of impacted third molars. *Acta Odontol Scand* 2002;60:108–12.
- [27] Billinton A. Human dental pulp as a source of native functional nociceptor pharmacology. *Pain* 2009;144:227.
- [28] Blatow M, Nennig E, Sarpaczki E, Reinhardt J, Schlieter M, Herweh C, Rasche D, Tronnier VM, Sartor K, Stippich C. Altered somatosensory processing in trigeminal neuralgia. *Hum Brain Mapp* 2009;30:3495–508.
- [29] Bongehiem U, Boissonade FM, Westermark A, Robinson PP, Fried K. Sympathetic nerve sprouting fails to occur in the trigeminal ganglion after peripheral nerve injury in the rat. *Pain* 1999;82:283–8.

- [30] Bonghenhielm U, Nosrat CA, Nosrat I, Eriksson J, Fjell J, Fried K. Expression of sodium channel SNS/PN3 and ankyrin(G) mRNAs in the trigeminal ganglion after inferior alveolar nerve injury in the rat. *Exp Neurol* 2000;164:384–95.
- [31] Bouloux GF. Temporomandibular joint pain and synovial fluid analysis: a review of the literature. *J Oral Maxillofac Surg* 2009;67:2497–504.
- [32] Bowles WR, Withrow JC, Lepinski AM, Hargreaves KM. Tissue levels of immunoreactive substance P are increased in patients with irreversible pulpitis. *J Endod* 2003;29:265–7.
- [33] Byers MR, Narhi MV. Dental injury models: experimental tools for understanding neuroinflammatory interactions and polymodal nociceptor functions. *Crit Rev Oral Biol Med* 1999;10:4–39.
- [34] Campbell RL, Parks KW, Dodds RN. Chronic facial pain associated with endodontic therapy. *Oral Surg Oral Med Oral Pathol* 1990;69:287–90.
- [35] Campbell WI, Kendrick RW. Pre-emptive analgesia using local anaesthesia: a study in bilaterally symmetrical surgery. *Br J Anaesth* 1997;79:657–9.
- [36] Carmichael FA, McGowan DA. Incidence of nerve damage following third molar removal: a West of Scotland Oral Surgery Research Group study. *Br J Oral Maxillofac Surg* 1992;30:78–82.
- [37] Cherkas PS, Huang TY, Pannicke T, Tal M, Reichenbach A, Hanani M. The effects of axotomy on neurons and satellite glial cells in mouse trigeminal ganglion. *Pain* 2004;110:290–8.
- [38] Cheshire Jr WP. Defining the role for gabapentin in the treatment of trigeminal neuralgia: a retrospective study. *J Pain* 2002;3:137–42.
- [39] Cheung LK, Lo J. The long-term clinical morbidity of mandibular step osteotomy. *Int J Adult Orthodon Orthognath Surg* 2002;17:283–90.
- [40] Chichorro JG, Zampronio AR, Cabrini DA, Franco CRC, Rae GA. Mechanisms operated by endothelin ETA and ETB receptors in the trigeminal ganglion contribute to orofacial thermal hyperalgesia induced by infraorbital nerve constriction in rats. *Neuropeptides* 2009;43:133–42.
- [41] Cittadini E, Matharu MS. Symptomatic trigeminal autonomic cephalalgias [Review]. *Neurologist* 2009;15:305–12.
- [42] Cook SP, Vulchanova L, Hargreaves KM, Elde R, McCleskey EW. Distinct ATP receptors on pain-sensing and stretch-sensing neurons. *Nature* 1997;387:505–8.
- [43] Cooper SA, Desjardins PJ. The value of the dental impaction pain model in drug development. *Methods Mol Biol* 2010;617:175–90.
- [44] Cruccu G, Biasiotta A, Di Rezzo S, Fiorelli M, Galeotti F, Innocenti P, Mameli S, Millefiorini E, Truini A. Trigeminal neuralgia and pain related to multiple sclerosis. *Pain* 2009;143:186–91.
- [45] Derry C, Derry S, Moore RA, McQuay HJ. Single dose oral ibuprofen for acute postoperative pain in adults. *Cochrane Database Syst Rev* 2009;CD001548.
- [46] Diatchenko L, Anderson AD, Slade GD, Fillingim RB, Shabalina SA, Higgins TJ, Sama S, Belfer I, Goldman D, Max MB, Weir BS, Maixner W. Three major haplotypes of the beta2 adrenergic receptor define psychological profile, blood pressure, and the risk for development of a common musculoskeletal pain disorder. *Am J Med Genet B Neuropsychiatr Genet* 2006;141B:449–62.
- [47] Diatchenko L, Slade GD, Nackley AG, Bhalang K, Sigurdsson A, Belfer I, Goldman D, Xu K, Shabalina SA, Shagin D, Max MB, Makarov SS, Maixner W. Genetic basis for individual variations in pain perception and the development of a chronic pain condition. *Hum Mol Genet* 2005;14:135–43.
- [48] Diogenes A, Akopian AN, Hargreaves KM. NGF up-regulates TRPA1: implications for orofacial pain. *J Dent Res* 2007;86:550–5.
- [49] Diogenes A, Patwardhan AM, Jeske NA, Ruparel NB, Goffin V, Akopian AN, Hargreaves KM. Prolactin modulates TRPV1 in female rat trigeminal sensory neurons. *J Neurosci* 2006;26:8126–36.
- [50] Dionne RA, Gordon SM, Rowan J, Kent A, Brahim JS. Dexamethasone suppresses peripheral prostanoid levels without analgesia in a clinical model of acute inflammation. *J Oral Maxillofac Surg* 2003;61:997–1003.
- [51] Dodick DW. Migraine with isolated facial pain: a diagnostic challenge. *Cephalalgia* 2007;27:1199–200.
- [52] Dolan JC, Lam DK, Achdjian SH, Schmidt BL. The dolognawmeter: a novel instrument and assay to quantify nociception in rodent models of orofacial pain. *J Neurosci Methods* 2010;187:207–15.
- [53] Dubner R, Bennett GJ. Spinal and trigeminal mechanisms of nociception. *Annu Rev Neurosci* 1983;6:381–418.
- [54] Dworkin SF, Huggins KH, LeResche L, Von Korff M, Howard J, Truelove E, Sommers E. Epidemiology of signs and symptoms in temporomandibular disorders: clinical signs in cases and controls. *J Am Dent Assoc* 1990;120:273–81.
- [55] Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord* 1992;6:301–55.
- [56] Dworkin SF, Sherman J, Mancl L, Ohrbach R, LeResche L, Truelove E. Reliability, validity, and clinical utility of the research diagnostic criteria for Temporomandibular Disorders Axis II Scales: depression, non-specific physical symptoms, and graded chronic pain. *J Orofac Pain* 2002;16:207–20.
- [57] Epstein JB, Hong C, Logan RM, Barasch A, Gordon SM, Oberlee-Edwards L, McGuire D, Napenas JJ, Elting LS, Spijkervet FKL, Brennan MT. A systematic review of orofacial pain in patients receiving cancer therapy. *Support Care Cancer* 2010;18:1023–31.
- [58] Eross E, Dodick D, Eross M. The Sinus, Allergy and Migraine Study (SAMS). *Headache* 2007;47:213–24.
- [59] Falace DA, Reid K, Rayens MK. The influence of deep (odontogenic) pain intensity, quality, and duration on the incidence and characteristics of referred orofacial pain. *J Orofac Pain* 1996;10:232–9.
- [60] Fehrenbacher JC, Sun XX, Locke EE, Henry MA, Hargreaves KM. Capsaicin-evoked iCGRP release from human dental pulp: a model system for the study of peripheral neuropeptide secretion in normal healthy tissue. *Pain* 2009;144:253–61.
- [61] Fillingim RB, Maixner W, Sigurdsson A, Kincaid S. Sexual and physical abuse history in subjects with temporomandibular disorders: relationship to clinical variables, pain sensitivity, and psychologic factors. *J Orofac Pain* 1997;11:48–57.
- [62] Flake NM, Bonebreak DB, Gold MS. Estrogen and inflammation increase the excitability of rat temporomandibular joint afferent neurons. *J Neurophysiol* 2005;93:1585–97.
- [63] Fried K, Bonghenhielm U, Boissonade FM, Robinson PP. Nerve injury-induced pain in the trigeminal system. *Neuroscientist* 2001;7:155–65.
- [64] Gaul C, Sandor PS, Galli U, Palla S, Ettlin DA. Orofacial migraine. *Cephalalgia* 2007;27:950–2.
- [65] Goncalves DA, Dal Fabbro AL, Campos JA, Bigal ME, Speciali JG. Symptoms of temporomandibular disorders in the population: an epidemiological study. *J Orofac Pain* 2010;24:270–8.
- [66] Gordon SM, Brahim JS, Dubner R, McCullagh LM, Sang C, Dionne RA. Attenuation of pain in a randomized trial by suppression of peripheral nociceptive activity in the immediate postoperative period. *Anesth Analg* 2010;95:1351–7.
- [67] Gordon SM, Brahim JS, Rowan J, Kent A, Dionne RA. Peripheral prostanoid levels and nonsteroidal anti-inflammatory drug analgesia: replicate clinical trials in a tissue injury model. *Clin Pharmacol Ther* 2002;72:175–83.
- [68] Gordon SM, Dionne RA, Brahim J, Jabir F, Dubner R. Blockade of peripheral neuronal barrage reduces postoperative pain. *Pain* 1997;70:209–15.
- [69] Goya T, Wakisaka S, Kinoshita K. Microvascular decompression for trigeminal neuralgia with special reference to delayed recurrence. *Neurol Med Chi (Tokyo)* 1990;30:462–7.
- [70] Gracely RH, Dubner R, Wolskee PJ, Deeter WR. Placebo and naloxone can alter post-surgical pain by separate mechanisms. *Nature* 1983;306:264–5.
- [71] Graham C, Bond SS, Gerkovich MM, Cook MR. Use of the McGill pain questionnaire in the assessment of cancer pain: replicability and consistency. *Pain* 1980;8:377–87.
- [72] Gregg JM. Neuropathic complications of mandibular implant surgery: review and case presentations. *Ann R Australas Coll Dent Surg* 2000;15:176–80.
- [73] Grushka M, Sessle BJ. Applicability of the McGill Pain Questionnaire to the differentiation of 'toothache' pain. *Pain* 1984;19:49–57.
- [74] Hamlyn PJ. Neurovascular relationships in the posterior cranial fossa, with special reference to trigeminal neuralgia. 1. Review of the literature and development of a new method of vascular injection-filling in cadaveric controls. *Clin Anat* 1997;10:371–9.
- [75] Hamlyn PJ. Neurovascular relationships in the posterior cranial fossa, with special reference to trigeminal neuralgia. 2. Neurovascular compression of the trigeminal nerve in cadaveric controls and patients with trigeminal neuralgia: quantification and influence of method. *Clin Anat* 1997;10:380–8.
- [76] Hamza M, Wang X-M, Adam A, Brahim JS, Rowan JS, Carmona GN, Dionne RA. Kinin B1 receptors contributes to acute pain following minor surgery in humans. *Mol Pain* 2010;6:12.
- [77] Hargreaves KM, Costello A. Glucocorticoids suppress levels of immunoreactive bradykinin in inflamed tissue as evaluated by microdialysis probes. *Clin Pharmacol Ther* 1990;48:168–78.
- [78] Hargreaves KM, Dionne RA, Mueller GP, Goldstein DS, Dubner R. Naloxone, fentanyl, and diazepam modify plasma beta-endorphin levels during surgery. *Clin Pharmacol Ther* 1986;40:165–71.
- [79] Harriott AM, Dessem D, Gold MS. Inflammation increases the excitability of masseter muscle afferents. *Neuroscience* 2006;141:433–42.
- [80] Harriott AM, Gold MS. Serotonin type 1D receptors (5HT_{1D}) are differentially distributed in nerve fibres innervating craniofacial tissues. *Cephalalgia* 2008;28:933–44.
- [81] Harriott AM, Gold MS. Contribution of primary afferent channels to neuropathic pain. *Curr Pain Headache Rep* 2009;13:197–207.
- [82] Harriott AM, Gold MS. Electrophysiological properties of dural afferents in the absence and presence of inflammatory mediators. *J Neurophysiol* 2009;101:3126–34.
- [83] Henry MA, Luo S, Foley BD, Rzasas RS, Johnson LR, Levinson SR. Sodium channel expression and localization at demyelinated sites in painful human dental pulp. *J Pain* 2009;10:750–8.
- [84] Hokfelt TS, Ceccatelli L, Gustafsson AL, Hulting V, Verge M, Villar X J, Xu ZQ, Xu Z, Wiesenfeld-Hallin, Zhang X. Plasticity of NO synthase expression in the nervous and endocrine systems. *Neuropharmacology* 1994;33:1221–7.
- [85] Hollins M, Harper D, Gallagher S, Owings EW, Lim PF, Miller V, Siddiqi MQ, Maixner W. Perceived intensity and unpleasantness of cutaneous and auditory stimuli: an evaluation of the generalized hypervigilance hypothesis. *Pain* 2009;141:215–21.
- [86] Hussain A, Stiles MA, Oshinsky ML. Pain remapping in migraine: a novel characteristic following trigeminal nerve injury. *Headache* 2010;50:669–71.
- [87] Inoue M, Fujii Y, Furukawa K, Okada M, Okumura K, Hayakawa T, Furukawa K, Sugiura Y. Refractory skin injury in complex knock-out mice expressing only the GM3 ganglioside. *J Biol Chem* 2002;277:29881–8.
- [88] Jannetta PJ. Neurovascular compression in cranial nerve and systemic disease. *Ann Surg* 1980;192:518–25.
- [89] Jeske NA, Patwardhan AM, Gamper N, Price TJ, Akopian AN, Hargreaves KM. Cannabinoid WIN 55, 212–2 regulates TRPV1 phosphorylation in sensory neurons. *J Biol Chem* 2006;281:32879–90.

- [90] Jimenez-Andrade JM, Mantyh WG, Bloom AP, Xu H, Ferng AS, Dussor G, Vanderah TW, Mantyh PW. A phenotypically restricted set of primary afferent nerve fibers innervate the bone versus skin: therapeutic opportunity for treating skeletal pain. *Bone* 2010;46:306–13.
- [91] John MT, LeResche L, Koepsell TD, Hujoel P, Miglioretti DL, Micheelis W. Oral health-related quality of life in Germany. *Eur J Oral Sci* 2003;111:483–91.
- [92] John MT, Miglioretti DL, LeResche L, Von Korff M, Critchlow CW. Widespread pain as a risk factor for dysfunctional temporomandibular disorder pain. *Pain* 2003;102:257–63.
- [93] Katusic S, Beard CM, Bergstralh E, Kurland LT. Incidence and clinical features of trigeminal neuralgia, Rochester, Minnesota, 1945–1984. *Ann Neurol* 1990;27:89–95.
- [94] Khan AA, Owatz CB, Schindler WG, Schwartz SA, Keiser K, Hargreaves KM. Measurement of mechanical allodynia and local anesthetic efficacy in patients with irreversible pulpitis and acute periradicular periodontitis. *J Endod* 2007;33:796–9.
- [95] Kim H, Lee H, Rowan J, Brahim J, Dionne RA. Genetic polymorphisms in monoamine neurotransmitter systems show only weak association with acute post-surgical pain in humans. *Mol Pain* 2006;2:24.
- [96] Kim H, Ramsay E, Lee H, Wahl S, Dionne RA. Genome-wide association study of acute post-surgical pain in humans. *Pharmacogenomics* 2009;10:171–9.
- [97] Kroner-Herwig B. Chronic pain syndromes and their treatment by psychological interventions. *Curr Opin Psychiatry* 2009;22:200–4.
- [98] Lee Y-S, Kim H, Wu T-X, Wang X-M, Dionne RA. Genetically mediated interindividual variation in analgesic responses to cyclooxygenase inhibitory drugs. *Clin Pharmacol Ther* 2006;79:407–18.
- [99] LeResche L. Epidemiology of temporomandibular disorders: implications for the investigation of etiologic factors. *Crit Rev Oral Biol Med* 1997;8:291–305.
- [100] LeResche L, Mancl LA, Drangsholt MT, Huang G, Von Korff M. Predictors of onset of facial pain and temporomandibular disorders in early adolescence. *Pain* 2007;129:269–78.
- [101] LeResche L, Mancl LA, Drangsholt MT, Saunders K, Korff MV. Relationship of pain and symptoms to pubertal development in adolescents. *Pain* 2005;118:201–9.
- [102] LeResche L, Mancl L, Sherman JJ, Gandara B, Dworkin SF. Changes in temporomandibular pain and other symptoms across the menstrual cycle. *Pain* 2003;106:253–61.
- [103] LeResche L, Saunders K, Von Korff MR, Barlow W, Dworkin SF. Use of exogenous hormones and risk of temporomandibular disorder pain. *Pain* 1997;69:153–60.
- [104] Levine JD, Gordon NC, Fields HL. The mechanism of placebo analgesia. *Lancet* 1978;2:654–7.
- [105] Levine JD, Gordon NC, Jones RT, Fields HL. The narcotic antagonist naloxone enhances clinical pain. *Nature* 1978;272:826–7.
- [106] Li YQ, Li H, Wei J, Qu L, Wu LA. Expression changes of K⁺-Cl⁻ co-transporter 2 and Na⁺-K⁺-Cl⁻ co-transporter1 in mouse trigeminal subnucleus caudalis following pulpal inflammation. *Brain Res Bull* 2010;81:561–4.
- [107] Lim PF, Smith S, Bhalang K, Slade GD, Maixner W. Development of temporomandibular disorders is associated with greater bodily pain experience. *Clin J Pain* 2010;26:116–20.
- [108] Lipton JA, Ship JA, Larach-Robinson D. Estimated prevalence and distribution of reported orofacial pain in the United States. *J Am Dent Assoc* 1994;124:115–21.
- [109] Lobb WK, Zakariassen KL, McGrath PJ. Endodontic treatment outcomes: do patients perceive problems? *J Am Dent Assoc* 1996;127:597–600.
- [110] Luo S, Perry GM, Levinson SR, Henry MA. Pulpitis increases the proportion of atypical nodes of Ranvier in human dental pulp axons without a change in Na v 1.6 sodium channel expression. *Neuroscience* 2010;169:1881–7.
- [111] Macfarlane TV, Blinkhorn AS, Craven R, Zakrzewska JM, Atkin P, Escudier MP, Rooney CA, Aggarwal V, Macfarlane GJ. Can one predict the likely specific orofacial pain syndrome from a self-completed questionnaire? *Pain* 2004;111:270–7.
- [112] Macfarlane TV, Blinkhorn AS, Davies RM, Ryan P, Worthington HV, Macfarlane GJ. Orofacial pain: just another chronic pain? Results from a population-based survey. *Pain* 2002;99:453–8.
- [113] Malinow K, Yannakakis GD, Glusman SM, Edlow DW, Griffin J, Pestronk A, Powell DL, Ramsey-Goldman R, Eidelman BH, Medsger TA, Jr. Subacute sensory neuropathy secondary to dorsal root ganglionitis in primary Sjögren's syndrome. *Ann Neurol* 1986;20:535–7.
- [114] Matthews B. The mechanisms of pain from dentine and pulp. *Br Dent J* 1976;140:57–60.
- [115] Mehlisch DR, Aspley S, Daniels SE, Southerden KA, Christensen KS. A single-tablet fixed-dose combination of racemic ibuprofen/paracetamol in the management of moderate to severe postoperative dental pain in adult and adolescent patients: a multicenter, two-stage, randomized, double-blind, parallel-group, placebo-controlled, factorial study. *Clin Ther* 2010;32:1033–49.
- [116] Melzack R. The McGill Pain Questionnaire: major properties and scoring methods. *Pain* 1975;1:277–99.
- [117] Melzack R, Jeans ME, Stratford JG, Monks RC. Ice massage and transcutaneous electrical stimulation: comparison of treatment for low-back pain. *Pain* 1980;9:209–17.
- [118] Miller JP, Acar F, Burchiel KJ. Classification of trigeminal neuralgia: clinical, therapeutic, and prognostic implications in a series of 144 patients undergoing microvascular decompression. *J Neurosurg* 2009;111:1231–4.
- [119] Morgan CR, Bird EV, Robinson PP, Boissonade FM. TRPA1 expression in human lingual nerve neuromas in patients with and without symptoms of dysesthesia. *Neurosci Lett* 2009;465:189–93.
- [120] Moskowitz MA, Macfarlane R. Neurovascular and molecular mechanisms in migraine headaches. *Cerebrovasc Brain Metab Rev* 1993;5:159–77.
- [121] Nackleley AG, Shabalina SA, Tchivileva IE, Satterfield K, Korchynskiy O, Makarov SS, Maixner W, Diatchenko L. Human catechol-O-methyltransferase haplotypes modulate protein expression by altering mRNA secondary structure. *Science* 2006;314:1930–3.
- [122] Nauta HJ, Wehman JC, Koliatsos VE, Terrell MA, Chung K. Intraventricular infusion of nerve growth factor as the cause of sympathetic fiber sprouting in sensory ganglia. *J Neurosurg* 1999;91:447–53.
- [123] Ni Riordain R, Moloney E, O'Sullivan K, McCreary C. Burning mouth syndrome and oral health-related quality of life: is there a change over time? *Oral Dis* 2010;16:643–7.
- [124] Oh IH, Choi SK, Park BJ, Kim TS, Rhee BA, Lim YJ. The treatment outcome of elderly patients with idiopathic trigeminal neuralgia: micro-vascular decompression versus gamma knife radiosurgery. *J Korean Neurosurg Soc* 2008;44:199–204.
- [125] Okada-Ogawa A, Suzuki I, Sessle BJ, Chiang C-Y, Salter MW, Dostrovsky JO, Tsuboi Y, Kondo M, Kitagawa J, Kobayashi A, Noma N, Imamura Y, Iwata K. Astroglia in medullary dorsal horn (trigeminal spinal subnucleus caudalis) are involved in trigeminal neuropathic pain mechanisms. *J Neurosci* 2009;29:11161–71.
- [126] Okamoto K, Bereiter DF, Thompson R, Tashiro A, Bereiter DA. Estradiol replacement modifies c-fos expression at the spinomedullary junction evoked by temporomandibular joint stimulation in ovariectomized female rats. *Neuroscience* 2008;156:729–36.
- [127] Owatz CB, Khan AA, Schindler WG, Schwartz SA, Keiser K, Hargreaves KM. The incidence of mechanical allodynia in patients with irreversible pulpitis. *J Endod* 2007;33:552–6.
- [128] Pajot J, Ressot C, Ngom I, Woda A. Gonadectomy induces site-specific differences in nociception in rats. *Pain* 2003;104:367–73.
- [129] Park S-S, Lee M-K, Kim J-W, Jung J-Y, Kim I-S, Ghang C-G. Percutaneous balloon compression of trigeminal ganglion for the treatment of idiopathic trigeminal neuralgia: experience in 50 patients. *J Korean Neurosurg Soc* 2008;43:186–9.
- [130] Patwardhan AM, Akopian AN, Ruparel NB, Diogenes A, Weintraub ST, Uhlson C, Murphy RC, Hargreaves KM. Heat generates oxidized linoleic acid metabolites that activate TRPV1 and produce pain in rodents. *J Clin Invest* 2010;120:1617–26.
- [131] Patwardhan AM, Berg KA, Akopian AN, Jeske NA, Gamper N, Clarke WP, Hargreaves KM. Bradykinin-induced functional competence and trafficking of the delta-opioid receptor in trigeminal nociceptors. *J Neurosci* 2005;25:8825–32.
- [132] Patwardhan AM, Diogenes A, Berg KA, Fehrenbacher JC, Clarke WP, Akopian AN, Hargreaves KM. PAR-2 agonists activate trigeminal nociceptors and induce functional competence in the delta opioid receptor. *Pain* 2006;125:114–24.
- [133] Patwardhan AM, Jeske NA, Price TJ, Gamper N, Akopian AN, Hargreaves KM. The cannabinoid WIN 55, 212–2 inhibits transient receptor potential vanilloid 1 (TRPV1) and evokes peripheral antihyperalgesia via calcineurin. *Proc Natl Acad Sci USA* 2006;103:11393–8.
- [134] Patwardhan AM, Scotland PE, Akopian AN, Hargreaves KM. Activation of TRPV1 in the spinal cord by oxidized linoleic acid metabolites contributes to inflammatory hyperalgesia. *Proc Natl Acad Sci USA* 2009;106:18820–4.
- [135] Pau AK, Croucher R, Marcenes W. Prevalence estimates and associated factors for dental pain: a review. *Oral Health Prev Dent* 2003;1:209–20.
- [136] Pepose JS, Foos RY, Stevens JG. Herpes simplex virus immediate-early protein ICP4 in murine models of latency. *Graefes Arch Clin Exp Ophthalmol* 1986;24:341–5.
- [137] Polycarpou N, Ng YL, Canavan D, Moles DR, Gulabivala K. Prevalence of persistent pain after endodontic treatment and factors affecting its occurrence in cases with complete radiographic healing. *Int Endod J* 2005;38:169–78.
- [138] Portenoy RK, Ugarte C, Fuller I, Haas G. Population-based survey of pain in the United States: differences among white, African American, and Hispanic subjects. *J Pain* 2004;5:317–28.
- [139] Price TJ, Helesic G, Parghi D, Hargreaves KM, Flores CM. The neuronal distribution of cannabinoid receptor type 1 in the trigeminal ganglion of the rat. *Neuroscience* 2003;120:155–62.
- [140] Putzki N, Pfriem A, Limmroth V, Yaldizli O, Tettgenborn B, Diener HC, Katsarava Z. Prevalence of migraine, tension-type headache and trigeminal neuralgia in multiple sclerosis. *Eur J Neurol* 2009;16:262–7.
- [141] Quang PN, Schmidt BL. Endothelin-A receptor antagonism attenuates carcinoma-induced pain through opioids in mice. *J Pain* 2010;11:663–71.
- [142] Robert C, Wilson CS, Donnadiou S, Gaudy JF, Arreto CD. Bibliometric analysis of the scientific literature on pain research: a 2006 study. *Pain* 2008;138:250–4.
- [143] Robinson PP, Smith KG. Lingual nerve damage during lower third molar removal: a comparison of two surgical methods. *Br Dent J* 1996;180:456–61.
- [144] Robinson JP, Willars GB, Tomlinson DR, Keen P. Axonal transport and tissue contents of substance P in rats with long-term streptozotocin-diabetes. Effects of the aldose reductase inhibitor 'statil'. *Brain Res* 1987;426:339–48.

- [145] Rowan MP, Ruparel NB, Patwardhan AM, Berg KA, Clarke WP, Hargreaves KM. Peripheral delta opioid receptors require priming for functional competence in vivo. *Eur J Pharmacol* 2009;602:283–7.
- [146] Ruparel NB, Patwardhan AM, Akopian AN, Hargreaves KM. Homologous and heterologous desensitization of capsaicin and mustard oil responses utilize different cellular pathways in nociceptors. *Pain* 2008;135:271–9.
- [147] Salas MM, Hargreaves KM, Akopian AN. TRPA1-mediated responses in trigeminal sensory neurons: interaction between TRPA1 and TRPV1. *Eur J Neurosci* 2009;29:1568–78.
- [148] Scaravilli F. Reduced substance P in hereditary sensory neuropathy in the mouse. *Brain Res* 1983;263:147–50.
- [149] Sessle BJ, Hu JW. Mechanisms of pain arising from articular tissues. *Can J Physiol Pharmacol* 1991;69:617–26.
- [150] Seymour RA, Charlton JE, Phillips ME. An evaluation of dental pain using visual analogue scales and the McGill Pain Questionnaire. *J Oral Maxillofac Surg* 1983;41:643–8.
- [151] Sigurdsson A, Maixner W. Effects of experimental and clinical noxious counterirritants on pain perception. *Pain* 1994;57:265–75.
- [152] Simpson MT, MacLaurin JG, Xu D, Ferguson KL, Vanderluit JL, Davoli MA, Roy S, Nicholson DW, Robertson GS, Park DS, Slack RS. Caspase 3 deficiency rescues peripheral nervous system defect in retinoblastoma nullizygous mice. *J Neurosci* 2001;21:7089–98.
- [153] Siqueira SR, Alves B, Malpartida HM, Teixeira MJ, Siqueira JT. Abnormal expression of voltage-gated sodium channels Nav1.7, Nav1.3 and Nav1.8 in trigeminal neuralgia. *Neuroscience* 2009;164:573–7.
- [154] Slade GD, Diatchenko L, Bhalang K, Sigurdsson A, Fillingim RB, Belfer I, Max MB, Goldman D, Maixner W. Influence of psychological factors on risk of temporomandibular disorders. *J Dent Res* 2007;86:1120–5.
- [155] Smeyne R J, Klein R, Schnapp A, Long LK, Bryant S, Lewin A, Lira SA, Barbacid M. Severe sensory and sympathetic neuropathies in mice carrying a disrupted Trk/NGF receptor gene. *Nature* 1994;368:246–9.
- [156] Smith TJ, Morrison LA, Leib DA. Pathogenesis of herpes simplex virus type 2 virion host shutoff (vhs) mutants. *J Virol* 2002;76:2054–61.
- [157] Spivack JG, Fareed MU, Valyi-Nagy T, Nash TC, O'Keefe JS, Gesser RM, McKie EA, MacLean AR, Fraser NW, Brown SM. Replication, establishment of latent infection, expression of the latency-associated transcripts and explant reactivation of herpes simplex virus type 1 gamma 34.5 mutants in a mouse eye model. *J Gen Virol* 1995;76:321–32.
- [158] Stephenson J, Byers MR. GFAP immunoreactivity in trigeminal ganglion satellite cells after tooth injury in rats. *Expt Neurol* 1995;131:11–22.
- [159] Serman AB, Schaumburg HH, Asbury AK. The acute sensory neuronopathy syndrome: a distinct clinical entity. *Ann Neurol* 1980;7:354–8.
- [160] Stohler CS, Kowalski CJ. Spatial and temporal summation of sensory and affective dimensions of deep somatic pain. *Pain* 1999;79:165–73.
- [161] Sweet WH. Deafferentation pain after posterior rhizotomy, trauma to a limb, and herpes zoster. *Neurosurgery* 1984;15:928–32.
- [162] Swift JQ, Garry MG, Roszkowski MT, Hargreaves KM. Effect of flurbiprofen on tissue levels of immunoreactive bradykinin and acute postoperative pain. *J Oral Maxillofac Surg* 1993;51:112–6 [discussion 116–7].
- [163] Tacconi L, Miles JB. Bilateral trigeminal neuralgia: a therapeutic dilemma. *Br J Neurosurg* 2000;14:33–9.
- [164] Tahmouh AJ. Causalgia: redefinition as a clinical pain syndrome. *Pain* 1981;10:187–97.
- [165] Tal M, Devor M. Ectopic discharge in injured nerves: comparison of trigeminal and somatic afferents. *Brain Res* 1992;579:148–51.
- [166] Tanaka T, Sakamoto E, Shiiba S, Oda M, Kito S, Wakasugi-Sato N, Matsumoto-Takeda S, Imamura Y, Nakanishi O, Morimoto Y. Relationship between the curative effects of carbamazepine administration and the neurovascular compression volume of the trigeminal nerve measured using magnetic resonance cisternography. *Clin J Pain* 2009;25:752–9.
- [167] Tarricone R, Aguzzi G, Musi F, Fariselli L, Casasco A. Cost-effectiveness analysis for trigeminal neuralgia: cyberknife vs microvascular decompression. *Neuropsychiatr Dis Treat* 2008;4:647–52.
- [168] Tashiro A, Okamoto K, Bereiter DA. Chronic inflammation and estradiol interact through MAPK activation to affect TMJ nociceptive processing by trigeminal caudalis neurons. *Neuroscience* 2009;164:1813–20.
- [169] Tchivileva IE, Lim PF, Smith SB, Slade GD, Diatchenko L, McLean SA, Maixner W. Effect of catechol-O-methyltransferase polymorphism on response to propranolol therapy in chronic musculoskeletal pain: a randomized, double-blind, placebo-controlled, crossover pilot study. *Pharmacogenet Genomics* 2010;20:239–48.
- [170] Thackray AM, Field HJ. Differential effects of famciclovir and valaciclovir on the pathogenesis of herpes simplex virus in a murine infection model including reactivation from latency. *J Infect Dis* 1996;173:291–9.
- [171] Tsuzuki K, Kondo E, Fukuoka T, Yi D, Tsujino H, Sakagami M, Noguchi K. Differential regulation of P2X(3) mRNA expression by peripheral nerve injury in intact and injured neurons in the rat sensory ganglia. *Pain* 2001;91:351–60.
- [172] U.S. Department of Health and Human Services. Oral health in America: a report of the Surgeon General. Rockville, MD: U.S. Department of Health and Human Services. National Institute of Dental and Craniofacial Research, National Institutes of Health; 2000.
- [173] Valmaseda-Castellon E, Berini-Ayres L, Gay-Escoda C. Lingual nerve damage after third lower molar surgical extraction. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000;90:567–73.
- [174] Vargas CM, Macek MD, Marcus SE. Sociodemographic correlates of tooth pain among adults: United States, 1989. *Pain* 2000;85:87–92.
- [175] Vaughn AH, Gold MS. Ionic mechanisms underlying inflammatory mediator-induced sensitization of dural afferents. *J Neurosci* 2010;30:7878–88.
- [176] Vickers ER, Cousins MJ, Woodhouse A. Pain description and severity of chronic orofacial pain conditions. *Aust Dent J* 1998;43:403–9.
- [177] Vos BP, Strassman AM, Maciewicz RJ. Behavioral evidence of trigeminal neuropathic pain following chronic constriction injury to the rat's infraorbital nerve. *J Neurosci* 1994;14:2708–23.
- [178] Warren CA, Mok L, Gordon S, Fouad AF, Gold MS. Quantification of neural protein in extirpated tooth pulp. *J Endod* 2008;34:7–10.
- [179] White TM, Gilden DH, Mahalingam R. An animal model of varicella virus infection. *Brain Pathol* 2001;11:475–9.
- [180] Wilkins RH. Neurovascular compression syndromes. *Neurol Clin* 1985;3:359–72.
- [181] Woodham P, Anderson P, Nadim W, Trueman M. Satellite cells surrounding axotomized rat dorsal root ganglion cells increase expression of GFAP-like protein. *Neurosci Lett* 1989;98:8–12.
- [182] Wu YW, Bi YP, Kou XX, Xu W, Ma LQ, Wang KW, Gan YH, Ma XC. 17-Beta-estradiol enhanced allodynia of inflammatory temporomandibular joint through upregulation of hippocampal TRPV1 in ovariectomized rats. *J Neurosci* 2010;30:8710–9.
- [183] Youle M, Osio M. A double-blind, parallel-group, placebo-controlled, multicentre study of acetyl L-carnitine in the symptomatic treatment of antiretroviral toxic neuropathy in patients with HIV-1 infection. *HIV Med* 2007;8:241–50.
- [184] Zakrzewska JM, Jassim S, Bulman JS. A prospective, longitudinal study on patients with trigeminal neuralgia who underwent radiofrequency thermocoagulation of the Gasserian ganglion. *Pain* 1999;79:51–8.
- [185] Zhang X, Ji RR, Arvidsson J, Lundberg JM, Bartfai T, Bedecs K, Hokfelt T. Expression of peptides, nitric oxide synthase and NPY receptor in trigeminal and nodose ganglia after nerve lesions. *Exp Brain Res* 1996;111:393–404.
- [186] Zhang JM, Song X, LaMotte R. Enhanced excitability of sensory neurons in rats with cutaneous hyperalgesia produced by chronic compression of the dorsal root ganglion. *J Neurophysiol* 2002;82:3359–66.